

A Systematic Review and Meta-Analysis About the Prevalence of Neck Pain in Fast Jet Pilots

Amelia Riches; Wayne Spratford; Jeremy Witchalls; Phil Newman

- BACKGROUND:** During flight, fast jet pilots frequently move their heads into extreme positions while withstanding large amounts of stress on their cervical spines. These factors are thought to contribute to episodes of neck pain.
- METHODS:** We conducted a systematic review and meta-analysis of previous neck pain prevalence data in fast jet pilots to determine an overall pooled prevalence. Subgroup analyses were performed according to when pilots complained about their neck pain, whether these same pilots sought treatment, and if they lost time from flying. Four research databases were searched. Studies were eligible for inclusion if they were written in English, involved a group of fast jet pilots who were actively flying high performance aircraft, and reported quantitative prevalence data about neck pain in these pilots. These eligibility criteria were independently applied by two reviewers and risk of bias was evaluated. MetaXL software was used to conduct the meta-analysis.
- RESULTS:** In total, 8003 fast jet pilots across 18 eligible studies were included in the review. The overall pooled prevalence of neck pain in fast jet pilots was 51%. It was found that 39% of subjects lost time from flying, while only 32% sought medical treatment.
- DISCUSSION:** Neck pain in fast jet pilots adversely affects operational capabilities of defense forces. The prevalence of neck pain varies according to the definitions or thresholds of complaints used across the literature. Further research is required to standardize the definition of neck pain.
- KEYWORDS:** frequency, cervical spine, fighter aircrew.

Riches A, Spratford W, Witchalls J, Newman P. A systemic review and meta-analysis about the prevalence of neck pain in fast jet pilots. *Aerosp Med Hum Perform.* 2019; 90(10):882–890.

The term ‘fast jet pilot’ describes military personnel who operate high performance aircraft.⁵¹ Their primary goal is to provide combat air power for their national government. This involves executing various airborne mission roles, some of which place high degrees of physical stress on the pilots.⁵¹ These pilots are also involved in planning missions, briefings, debriefings, and other administrative tasks which are completed in office settings.^{11,51}

A skill required for aerial combat is the ability to move the aircraft while executing offensive and defensive flight maneuvers. This is called air combat maneuvering, and is essential for the pilot to perform optimally within a combat environment.^{39,42} Throughout these tactical movements, fast jet pilots constantly turn their heads to scan for targets in the air and on the ground.²⁹ Hence, air combat maneuvering involves rapid and repetitive excursions of the cervical spine away from the anatomically neutral position.^{29,54,55} Fast jet pilots complete these maneuvers while they are also exposed to high gravitational

accelerations (G force or G),^{28,30} which increase compression and stress along the axis of the spine.^{30,39,52} Axial loads equivalent to 65 kg are exerted on the neck due to the weight of the head, helmet and helmet-mounted equipment when pilots fly at 9 G.^{2,26} Although some cross sectional studies suggest that exposure to these forces can initially strengthen the neck,^{1,18} it is widely acknowledged that they likely contribute to acute and/or chronic episodes of neck pain.^{1,37,51} Due to these physically demanding occupational factors, it is expected that fast jet pilots will have a higher prevalence of neck pain

From The Faculty of Health, Discipline of Physiotherapy at University of Canberra, Canberra, Australia.

This manuscript was received for publication in February 2019. It was accepted for publication in July 2019.

Address correspondence to: Amelia Riches, Locked Bag 1, University of Canberra, ACT 2601 Australia; amelia_riches@live.com.au.

Reprint & Copyright © by the Aerospace Medical Association, Alexandria, VA.

DOI: <https://doi.org/10.3357/AMHP.5360.2019>

compared to that of the general population.²⁴ The prevalence of neck pain within the global community is 4.9%, where 90% of cases are attributed to injuries of the intervertebral discs, 80% to anterior spinal ligaments, and 40% to facet joints following vehicle collisions.⁹ Neck pain can result in lost workdays and reduced flying capacities for fast jet pilots.^{37,51} Given that \$15.2 million is required to train one pilot for an F/A-18 Hornet aircraft,³⁶ days lost from flying are expensive to defense forces⁴⁸ and informed risk strategies are required. This issue has gained international focus and is supported by a specialized North Atlantic Treaty Organization (NATO) Research Task Group: HFM 252.⁴³

The prevalence of neck pain in fast jet pilots must be determined to understand the magnitude of the issue and the characteristics of those fast jet pilots who are affected. This is important due to the growing advances in aircraft technology and helmet-mounted devices, which will influence the forces applied to the neck.² To date, there is only one published systematic review and meta-analysis which examines neck pain in the fast jet pilot population.⁴⁵ However, this study did not define neck pain⁴⁵ which reduced the clarity of research findings.

There is large variation in how pain and injuries are reported in epidemiology research.⁵⁰ They typically fall into one of three definitions; 'all complaints reported,' 'time lost,' and 'medical attention sought'.^{8,50} 'All complaints reported' refers to the expression of any health-related incident irrespective of its ramifications or the interventions required.⁸ Injuries defined by 'time lost' are those which impair an individual's ability to fully engage in training or activity, while 'medical attention sought' refers to injuries requiring medical assessment.^{8,50} With several definitions of neck pain and injury available to researchers, there is little comparable prevalence data which makes drawing clear conclusions from the literature challenging. The present review aims to address this gap and clearly convey the prevalence of neck pain in fast jet pilots. The focused questions for this review are:

1. What is the prevalence of neck pain in fast jet pilots?
2. How does prevalence vary across four thresholds of complaints (i.e., 'any complaint experienced at any time,' 'any complaint experienced during a specified time,' 'medical attention sought,' and 'time lost from flying')?

METHODS

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were followed when completing this review.³¹ The protocol for the review was registered on the 19th of February 2018 in the PROSPERO international prospective register for systematic reviews website (<https://www.crd.york.ac.uk/prospero/>), registration number CRD42018088909. The review did not meet the criteria outlined by the University of Canberra's Human Research Ethics Committee and therefore did not require ethical approval.

Procedure

Four electronic databases including Medline, CINAHL, Scopus, and Google Scholar were searched for literature published

between the years 2000 and 2018. Search terms related to fast jet pilots, prevalence, and neck pain. Searches were repeated to identify papers published between review completion and submission in February 2019.

Titles and abstracts were screened separately by two reviewers to identify relevant studies. Both reviewers retrieved full text copies of these papers and independently examined them for relevance. Reference lists of each study were scanned (forward and backward citation tracking) to ensure all eligible papers were identified. Blinding throughout was achieved by using the Covidence software package.¹⁰ Disagreement between the reviewers was resolved thereafter by consensus.

Predetermined eligibility criteria were used to screen titles, abstracts, and full texts. Studies were eligible for inclusion if they met these criteria: 1) were written in English; 2) involved a discernible group of fast jet pilots who were actively flying; and 3) reported quantitative data about the prevalence of neck pain in these pilots.

Studies were excluded for the following reasons: 1) they were not written in English; 2) included fast jet pilots who could not be easily distinguished from a larger sample of pilots; or 3) were case studies, expert opinions or reviews.

The literature considers aircraft capable of flying at 5 G or more high performance 'airframes'.^{1,2,54} Since the majority of subjects operated high performance aircraft, their exposure to increased G force was high. The reviewers agreed that operating high performance aircraft and therefore flying at high G force was the exposure of interest.

The number of fast jet pilots, age, sex, military flying hours, and type of aircraft flown were recorded to determine the parity and differences between studies (**Table I**).

The outcome of interest was the prevalence of neck pain in fast jet pilots. Prevalence was recorded as the number of fast jet pilots who reported neck pain out of the total number of fast jet pilots sampled (Table I). When a prevalence value was not specified, a proportion was calculated when and where data presented in the papers allowed. Prevalence was documented according to four thresholds for complaints including 'any complaint experienced at any time,' 'any complaint experienced at a specified time,' 'time lost from flying,' and 'medical treatment sought.' Only the proportions of affected pilots who flew high performance aircraft were recorded for analysis.

Selection, nonresponse and measurement biases of the included studies were assessed independently by two researchers using the Quality assessment Checklist for Prevalence Studies.²³ The tool was designed for the evaluation of observational prevalence studies. It contains nine dichotomous items which assess external and internal validity while providing a summary score.²³ Literature shows that the overall interrater agreement of the tool is 91% with a 'strong' Kappa agreement of 0.82.^{23,34} Therefore, the instrument was considered appropriate given its high reliability and applicability to prevalence research.

Statistical Analysis

Data were extracted from the studies and cross-checked by a second reviewer. The methods of each paper including the

Table I. Summary of the Included Studies.

PAPER	SUBJECTS	AIRCRAFT	DESIGN	DEFINITION OF NECK PAIN	PREVALENCE
Alricsson, 2004 ¹	N = 40 Age: - RG: 29.4 (± 4.5) - NRG: 29.4 (± 3.1)	Not specified	Cross-sectional	Not specified	AST: 17/40 (42.5%)
Ang, 2005 ²	N = 60 Age: 37 (± 7.5)	JAS 39 Gripen and JA 37 Viggen	Cross-sectional	"... defined as any neck-pain experience, neck stiffness or tenderness."	AST: 16/30 (53.3%)
De Loose, 2008 ¹¹	N = 90 Age: not specified	F-16	Cross-sectional	"... more than two episodes of neck pain [lasting] at least 1 [day during] the past 12 [months]... defined as pain in the head and neck..."	AST: 17/90 (18.9%) ST: 4/17 (23.5%) FTL: 6/17 (35.3%)
Drew, 2000 ¹³	N = 79 Age: 31.4 (± 4.5)	F-16 and F-15	Cross-sectional	"... specific spinal symptoms which occurred... [and their] effects... on aviator flying performance..."	AAT: 19/35 (54.3%) AST: 12.6/35 (36.0%) ST: 15/19 (78.9%) FTL: 6/19 (31.6%)
Greeves, 2006 ¹⁶	N = 144 Age: not specified	Not specified	Cross-sectional	Not specified	AAT: 100/144 (69.9%) AST: 23/144 (16.0%) ST: 31/100 (31.0%) AAT: 101/214 (47.2%)
Grossman, 2012 ¹⁸	N = 566 Age: 31.21 (± 8.15)	F-15, F-16, Skyhawk	Cross-sectional	"... [any back pain]... recalled. Location... was categorized as cervical,... thoracic,... and... lumbar... pain"	ST: 34/5681 (0.6%)
Hermes, 2010 ²⁰	N = 19673 Age: - DOB < 1966: 6279 (± 31.9) - DOB > 1967: 13394 (± 68.1)	Not specified.	Cross-sectional	"... presence of an initial waiver in the [aeromedical] database where the primary diagnosis was related to a cervical or lumbar intervertebral disc bulge, disc herniation or protrusion, disc degeneration, and/or spondylosis."	AAT: 25/27 (92.6%)
Jones, 2000 ²⁶	N = 70 Age: not specified.	C-26, KC-135, G-2, T-38, F-15, F-16 and F/A-18	Cross-sectional	"The questionnaire... sought to evaluate the presence, nature, severity, and timing of symptoms..."	AAT: 483/998 (48.4%)
Kang, 2011 ²⁷	N = 1003 Age: 29.3 (± 5.1)	Fighters: F-4, F-5, F-15, F-16 Trainers: KT-1, T-50, T-59	Cross-sectional	"... neck pain was defined as the presence of any neck pain or pain radiating to an upper extremity..."	AST: 0/10 (0%)
Landau, 2006 ²⁸	N = 30 Age: 35.2 (± 4)	F-4, F-15 and F-16	Cross-sectional	"... the duration and frequency of off-flight symptoms on average retrospectively per year"	AAT: 56/58 (96.6%) AST: 48/58 (82.8%) ST: 29/56 (51.8%) FTL: 24/56 (42.9%)
Lange, 2011 ²⁹	N = 58 Age: not specified.	F-16	Cross-sectional	"[the questionnaire] asked for the frequency of any pain... in the spine... [subjects] were asked to identify area(s) of discomfort on a figure of the human back"	AST: 27/32 (84.4%)
Moon 2015 ³⁵	N = 63 Age: - No neck pain group: 30.6 (± 3.88) - Neck pain group: 29.4 (± 3.85)	F-4, F-5, F-16, F-15, and T-50	Cross-sectional	"... neck pain during the past 1 yr... frequency, duration, and referred pain were investigated..."	AAT: 78/82 (95.1%)
Netto 2011 ³⁷	N = 82 Age: 31 (± 7.4)	Hawk 127 and F/A-18	Cross-sectional	"... the survey assessed neck strain, pain or injury..."	AST: 69/113 (61.1%)
Rintala 2015 ⁴²	N = 195 Age: 30 (± 5)	Not specified.	Cross-sectional	"The location of flight-induced musculoskeletal symptoms over the past 6 months was marked in each subject's pain observation chart"	

Table 1, Continued

PAPER	SUBJECTS	AIRCRAFT	DESIGN	DEFINITION OF NECK PAIN	PREVALENCE
Thoolen, 2015 ⁴⁹	N = 59 Age: - < 30 yr: 29% - 30-40 yr: 46% - > 40 yr: 25%	F-16	Cross-sectional	"Pain was defined as any pain or discomfort... A figure was used to define... the neck and lower back."	AAT: 35/59 (59.3%)
VanValkenburg, 2016 ⁵⁴	N = 250 Age: - SP's 20-29: 90% - IP's 25-34: 70% - IP's 35-44: 24%	T-6, T-38 and T-1 SUPT	Cross-sectional	Not specified	AAT: 92/250 (36.8%)
Verde, 2015 ⁵⁵	N = 70 Age: - F-16: 32.54 (± 4.71) - Typhoon: 33.37 (± 3.84)	F-16 and Eurofighter Typhoon	Cross-sectional	"...at least two episodes of neck pain after flight in the last 2 months."	AAT: 19/35 (54.3%)
Wagstaff, 2012 ⁵⁶	N = 105 Age: - 20-29: 47 - 30-39: 45 - 40-45: 7 - > 50: 6	F-16, F-14, F-5, T-38, T-37, and CF-104	Cross-sectional	"Neck pain events in general and specifically in the last 12 months."	AAT: 76/105 (72.4%) AST: 53/105 (50.5%)

design, definition of neck pain used and outcome data (i.e., prevalence of neck pain) were recorded (Table 1). Prevalence values from all studies were then subject to meta-analysis to determine a pooled prevalence of neck pain. This was conducted using MetaXL software version 5.3.⁶ In accordance with published recommendations,⁵ a double arcsine transformation was applied across the data to limit the overweighting associated with inverse variance analysis of pooled effects.⁵ Since the I^2 statistic exceeded 50%, a random effects model was used. Subgroup analyses were then performed according to the four thresholds of complaint, where the proportions of pilots who sought medical treatment and lost time from flying were determined from the group of pilots with neck pain. Subgroup analyses were completed if there were two or more comparable studies.

RESULTS

A total of 176 potentially relevant studies were returned from the database searches (Fig. 1). After duplicates were removed, 125 papers remained. Titles and abstracts were independently screened against the eligibility criteria, with 83% agreement between the two reviewers. Once disagreements were resolved by discussion, 30 papers were left for full text appraisal. The level of agreement between the reviewers during full text evaluation was 93% with disputes being resolved by consensus. Eighteen studies met the eligibility criteria and were included for review.

A total of 8003 fast jet pilots were anonymously surveyed about neck pain across the 18 studies. The mean age of the subjects ranged from 29.3 to 37 yr (Table 1). After the studies were categorized under the thresholds for complaints, 11 were found to define neck pain as 'any complaint experienced at any time' while nine described it as 'any complaint experienced during a specified time' (in the previous 3, 6, or 12 mo) and three reported it as both. Five studies defined neck pain by 'medical treatment sought,' while three of these also established their threshold as 'time lost from flying.'

The fast jet pilots included in 11 of the reviewed studies operated the F-16 Fighting Falcon (Table 1).

Twelve studies directly reported the prevalence of neck pain as proportions. We therefore calculated prevalence values from data presented in the remaining six papers (Table 1). This standardized the results to enable data combination for the meta-analysis and facilitated easier comparisons between studies.

According to the guidelines for classifying summary scores in the quality assessment tool, the studies received a mean score of 2.47 out of 9 (± 1.37) indicating a low risk of bias.²³ The reliability and validity of the questionnaires used for data collection were not reported or were unknown in all papers except three, and were consequently rated at high risk of bias (Fig. 2). Eight studies scored poorly in criteria associated with nonresponse bias. In contrast, each paper scored well for criteria eight, demonstrating that the mode for data collection was consistent within each study.




Fig. 1. PRISMA flow chart for study selection.

Interrater reliability for the quality assessment was calculated using the first-order agreement coefficient (AC1).¹⁹ Unlike Cohen's Kappa, the AC1 is not skewed by trait prevalence or marginal probabilities and offers a more stable reflection of interrater reliability.^{19,58} The AC1 for the quality assessment was calculated as 0.34, indicating that the level of agreement between both reviewers was 'fair' when assessing for risk of bias.

After pooling the prevalence data, a meta-analysis was performed (Fig. 3). The I^2 statistic was calculated at 99% indicating considerable heterogeneity between the studies.²¹ Since an adequate number of comparable studies were found, subgroup analyses were conducted according to each threshold for complaint.

The pooled prevalence of neck pain in fast jet pilots was 51% (95% CI, 33–68%) (Fig. 3). The prevalence of neck pain defined as 'any complaint experienced at any time,' was 71% (95% CI, 56–83%). It was found that 46% (95% CI, 0.31–0.61%) of fast jet pilots reported neck pain described as 'any complaint experienced during a specified time,' (3 to 12 mo). Of the fast jet pilots who reported neck pain, 31% (95% CI, 0–75%) sought treatment while 39% (95% CI, 30–50%) reported time lost from flying.

DISCUSSION

This review demonstrates that there is a high (51%) prevalence of neck pain among fast jet pilots. The sub-grouped analyses show that prevalence can be influenced by the definitions of neck pain or thresholds for complaints used in the research. This study therefore explains the reasons for the high variability of neck pain prevalence in fast jet pilots across the literature.

The meta-analysis demonstrates that only 32% of fast jet pilots with neck pain sought medical attention, while 39% lost time from flying. Further studies are required to explore why

medical attention is not commonly sought by fast jet pilots who experience neck pain and to encourage those affected personnel to seek treatment. Additional research is also warranted to investigate solutions which prevent the initial onset of neck pain in this population.

The studies included in this review demonstrated a considerable level of heterogeneity ($I^2 = 99\%$). This may relate to the methodological diversity of the studies, where neck pain was inconsistently defined.²⁵ For example, some researchers defined neck pain as symptoms which occurred during the last 3, 6, or 12 mo. However, other studies described neck pain as any neck symptom which occurred at any time over previous years. Therefore, inconsistent

definitions of neck pain have likely affected the results across the papers.

The studies also provided variable descriptions of the neck. Some papers defined this area by providing shaded sections on body charts, while others encouraged subjects to mark where they had experienced neck pain on body diagrams. Pooling data from research founded on non-standardized definitions leads to high variation in the collection and assessment of results.⁴¹ However, robust statistical adjustments⁵ were applied to control for this heterogeneity and improve the generalizability of findings. Therefore, the use of non-standardized definitions has been identified as a weakness in the literature.^{27,35} This gap may be addressed through Delphi studies, which will likely produce uniform definitions of both neck pain and neck regions to improve the accuracy of future results.

The risk of bias for the papers was possibly influenced by the lack of demonstrated validity and reliability of questionnaires used across the research. Although it is recommended that studies disclose the validity and reliability of their methods for data collection,⁴⁴ 15 of the included studies did not report these measures. However three papers modeled their surveys from previously validated questionnaires like the Dutch musculoskeletal questionnaire and the neck disability index. Since these tools are standardized and valid instruments,^{22,33} only these studies were assessed as having a low risk of bias for the seventh domain of the quality assessment.²³ If the questionnaires used in the 15 remaining studies were of poor validity and reliability, then a higher degree of random error may have affected the results⁵⁷ and masked the prevalence of neck pain.

It is expected that low participant response rates in eight of the included studies also affected the risk of bias and pooled prevalence. Nonresponse bias is a common limitation in studies which collect data through surveys or questionnaires.^{7,12} As

	Hoy 1	Hoy 2	Hoy 3	Hoy 4	Hoy 5	Hoy 6	Hoy 7	Hoy 8	Hoy 9	Hoy 10
Alicsson 2004	+	+	-	-	+	-	-	+	+	?
Ang 2005	-	+	+	+	+	+	-	+	+	+
DeLoose 2008	+	+	-	-	+	+	+	+	+	+
Drew 2000	+	+	+	-	+	+	-	+	+	+
Greeves 2006	+	+	+	-	+	-	+	+	+	+
Grossman 2012	+	+	+	+	+	+	-	+	+	+
Hermes 2010	+	+	+	-	-	+	-	+	+	+
Jones 2000	-	+	+	-	+	-	-	+	+	?
Kang 2011	+	+	+	+	+	+	-	+	+	+
Landau 2006	-	-	-	+	+	-	-	+	-	?
Lange 2011	+	+	+	+	+	+	-	+	+	+
Moon 2015	+	+	+	+	-	+	-	+	+	+
Netto 2011	+	+	+	+	+	-	-	+	+	+
Rintala 2015	+	+	+	+	+	+	-	+	+	+
Thoolen 2015	+	+	+	-	+	+	+	+	+	+
VanValkenburg 2016	+	+	+	+	+	-	-	+	-	+
Verde 2015	+	+	+	+	+	+	-	+	+	+
Wagstaff 2012	+	+	+	-	+	+	-	+	-	+

Fig. 2. Risk of bias scores for the included studies ($N = 18$). Legend: Hoy 1 = representative population; Hoy 2 = appropriate sampling frame; Hoy 3 = random selection or census; Hoy 4 = non-response bias; Hoy 5 = data collection; Hoy 6 = case defined; Hoy 7 = reliability and validity; Hoy 8 = direct data collection; Hoy 9 = numerators and denominators; Hoy 10 = summary risk of bias. Light grey (+) = 'low' risk of bias; light grey (?) = 'moderate' risk of bias; dark grey (-) = 'high' risk of bias.

acknowledged by several studies evaluated throughout this paper, any lack of total anonymity negatively influences voluntary response rates in military research.^{26,47,56} Although the included studies de-identified subjects, some pilots may not have answered the questionnaires due to risk of recognition by variables like age or flying hours.⁴⁷ Literature has shown that responders to health surveys are more likely to be well, and therefore report better health statuses compared to non-responders.^{7,40,53} Hence, the prevalence of neck pain has likely been underestimated throughout the papers which may have influenced the results of the meta-analysis.

ity between subjects in this review. It should be noted that several papers investigated their respective national populations of fast jet pilots, which may also explain similarities found across the subjects.^{2,27,38} Therefore it is likely that the pilots sampled in each study were representative of their national fast jet pilot populations, which may have enhanced the generalizability of the results.

The F-16 Fighting Falcon was the most common aircraft operated by fast jet pilots across the studies. However, several papers sampled fast jet pilots who flew more diverse ranges of high performance airframes (e.g., student versus instructor

All studies except one collected data through surveys or questionnaires, indicating that the mode of data collection was consistent across the majority of papers. Only one study obtained records from an electronic aeromedical database.²⁰ Results from an ongoing prospective cohort study of active military personnel¹⁵ have shown a moderate level of agreement between information gained from medical records and questionnaires.⁴⁶ Collecting data from medical records produces more statistically powerful, homogeneous and generalizable results and is therefore the gold standard for data collection in epidemiological studies.^{3,17} Nevertheless, this method is not immune from bias due to the typically limited disclosure of medical symptoms within military populations.^{7,20,56} Although the larger statistical power of this study may have affected the heterogeneity between the papers, it is clear that a consistent approach was used to collect data across the reviewed literature.

The studies sampled fast jet pilots who shared similarities in relation to gender, age and flying hours. Hence, the pilots exhibited relatively homogenous participant characteristics. Comprehensive testing procedures are used to select candidates who are best suited for operating high performance aircraft,³⁸ and involve evaluating cognitive, spatial, perceptual and psychomotor abilities as well as anthropometry.¹⁴ These standardized assessments are likely to explain the homogeneity




Fig. 3. Forest plot of pooled prevalence and subgroup analyses for neck pain in fast jet pilots.

aircraft). The forces imposed on fast jet pilots depend on the type of aircraft flown.⁴ None of the studies included in this review specified the proportions of fast jet pilots operating each type of airframe. Although a subgrouped meta-analysis according to types of aircraft could not be performed, the combination of data from pilots of different aircraft may have enhanced the generalizability of the results to a wider population.

Agreement between the reviewers was 83% during title and abstract screening and 93% during full text evaluation. Although there are no established standards which define 'satisfactory' agreement, literature indicates that consensus levels exceeding 80% are generally acceptable.³² However, when the studies were evaluated for risk of bias with the quality assessment tool, the AC1 statistic of 0.34 indicated a 'fair' agreement between the

two reviewers.^{19,23} It is our impression that the dichotomous nature of the instrument contributed to this level of consensus. Had the tool allowed for three choices during assessment (high, low and unclear), the reliability between the reviewers may have improved. Additionally, consensus may have been enhanced if the reviewers became more familiar with the tool through a training period. Agreement may have also improved if the reviewers standardized the interpretations of the domains with each other prior to assessing the papers.

A number of limitations must be considered when interpreting the results from this review. All studies failed to report the degree of neck pain (mild, moderate, or severe) and the proportions of fast jet pilots operating each type of aircraft. This limited our ability to investigate prevalence according to pain severity and types of airframes. Therefore our results only provide information about the prevalence of neck pain according to four thresholds for complaints. Despite these limitations, this review effectively synthesized literature applicable to the research questions, identified gaps in knowledge and suggested directions for future study.

This review calculated the pooled prevalence of neck pain in fast jet pilots as 51%, which is considerably higher than that of the global population. Several gaps in knowledge and literature were identified, including inconsistencies in defining neck pain and neck regions. This review demonstrated a limited use of validated and reliable questionnaires as well as high degrees of nonresponse bias across the research. It is recommended that subsequent researchers consider these limitations and regulate the approach for defining and assessing neck pain in fast jet pilots. Once achieved, the overall precision of results may increase, thereby allowing for more accurate conclusions. Future research should aim to establish the global prevalence of neck pain in fast jet pilots using internationally standardized methods of assessment. The mechanisms which cause neck pain and injury should also be investigated to form the basis of risk management and treatment programs.

ACKNOWLEDGMENTS

We would like to thank Dr. Angie Fearon for advice regarding the ACI statistic.

Financial Disclosure Statement: The authors had no competing interests to declare.

Funding: This review did not receive funding.

Authors and affiliations: Amelia Riches, B. Physiotherapy (Hons.), Wayne Spratford, Ph.D., Jeremy Witchalls, Ph.D., and Phil Newman, Ph.D., The Faculty of Health, Discipline of Physiotherapy at University of Canberra, Australia.

REFERENCES

1. Alricsson M, Harms-Ringdahl K, Larsson B, Linder J, Werner S. Neck muscle strength and endurance in fighter pilots: effects of a supervised training program. *Aviat Space Environ Med.* 2004; 75(1):23–28.
2. Äng B, Linder J, Harms-Ringdahl K. Neck strength and myoelectric fatigue in fighter and helicopter pilots with a history of neck pain. *Aviat Space Environ Med.* 2005; 76(4):375–380.
3. Argyrous G. *Statistics for research: with a guide to SPSS.* Singapore: SAGE Publications; 2011:509–511.
4. Arnott EJ. *A new beginning in sight.* London (UK): Hodder Arnold; 2006:141.
5. Barendregt JJ, Doi SA, Lee YY, Norman RE, Vos T. Meta-analysis of prevalence. *J Epidemiol Community Health.* 2013; 67(11):974–978.
6. Barendregt JJ, Doi SA. *MetaXL user guide Version 5.3.* MetaXL publications; 2016:1–85.
7. Cheung KL, Peter M, Smit C, de Vries H, Pieterse ME. The impact of non-response bias due to sampling in public health studies: a comparison of voluntary versus mandatory recruitment in a Dutch national survey on adolescent health. *BMC Public Health.* 2017; 17(1):276.
8. Clarsen B, Bahr R. Matching the choice of injury/illness definition to study setting, purpose and design: one size does not fit all! *Br J Sports Med.* 2014; 48(7):510–512.
9. Clemens HJ, Burow K. Experimental investigation on injury mechanisms of cervical spine at frontal and rear-front vehicle impacts. *SAE Trans.* 1972; 81:2779–2796.
10. Covidence. *Better systematic review management 2018* [Internet]. [Accessed 01 Aug. 2019.] Available from: <https://www.covidence.org/home>.
11. De Loose V, Van den Oord M, Burnotte F, Van Tiggelen D, Stevens V, et al. Individual, work-, and flight-related issues in F-16 pilots reporting neck pain. *Aviat Space Environ Med.* 2008; 79(8):779–783.
12. Dillman DA. *Mail and telephone surveys: the total design method.* New York (NY): John Wiley & Sons; 2007:89–98.
13. Drew WE, Sr. Spinal symptoms in aviators and their relationship to G-exposure and aircraft seating angle. *Aviat Space Environ Med.* 2000; 71(1):22–30.
14. Gradwell D, Rainford DJ, editors. *Ernsting's aviation medicine, 4th ed.* London: CRC Press; 2006: 309–310.
15. Gray GC, Chesbrough KB, Ryan MA, Amoroso P, Boyko EJ, et al. The Millennium Cohort Study: a 21-year prospective cohort study of 140,000 military personnel. *Mil Med.* 2002; 167(6):483–488.
16. Greeves J, Wickes MS. Review of the United Kingdom national work programme on the long term effects of sustained high G on the cervical spine. Neuilly-sur-Seine (France): NATO Research and Technology Organisation; 2014. Report No.: AC/323(HFM-083)TP/205.
17. Gregory KE, Radovinsky L. Research strategies that result in optimal data collection from the patient medical record. *Appl Nurs Res.* 2012; 25(2):108–116.
18. Grossman A, Nakdimon I, Chapnik L, Levy Y. Back symptoms in aviators flying different aircraft. *Aviat Space Environ Med.* 2012; 83(7): 702–705.
19. Gwet K. Inter-rater reliability: dependency on trait prevalence and marginal homogeneity. *Statistical Methods for Inter-Rater Reliability Assessment Series.* 2002; 2:1–9.
20. Hermes ED, Webb TS, Wells TS. Aircraft type and other risk factors for spinal disorders: data from 19,673 military cockpit aircrew. *Aviat Space Environ Med.* 2010; 81(9):850–856.
21. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ.* 2003; 327(7414):557–560.
22. Hildebrandt VH, Bongers P, Van Dijk F, Kemper HC, Dul J. Dutch Musculoskeletal Questionnaire: description and basic qualities. *Ergonomics.* 2001; 44(12):1038–1055.
23. Hoy D, Brooks P, Woolf A, Blyth F, March L, et al. Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. *J Clin Epidemiol.* 2012; 65(9):934–939.
24. Hoy D, March L, Woolf A, Blyth F, Brooks P, et al. The global burden of neck pain: estimates from the Global Burden of Disease 2010 study. *Ann Rheum Dis.* 2014; 73(7):1309–1315.
25. Jackson S. *Research Methods: A Modular Approach.* Independence (KY): Cengage Learning; 2007:57–60.
26. Jones JA, Hart SF, Baskin DS, Effenhauser R, Johnson SL, et al. Human and behavioral factors contributing to spine-based neurological cockpit injuries in pilots of high-performance aircraft: recommendations for management and prevention. *Mil Med.* 2000; 165(1):6–12.

27. Kang S, Hwang S, Lee ET, Yang S, Park J. Measuring the cumulative effect of G force on aviator neck pain. *Aviat Space Environ Med.* 2011; 82(11):1042–1048.
28. Landau DA, Chapnick L, Yoffe N, Azaria B, Goldstein L, Atar E. Cervical and lumbar MRI findings in aviators as a function of aircraft type. *Aviat Space Environ Med.* 2006; 77(11):1158–1161.
29. Lange B, Torp-Svendsen J, Toft P. Neck pain among fighter pilots after the introduction of the JHMCS helmet and NVG in their environment. *Aviat Space Environ Med.* 2011; 82(5):559–563.
30. Lecompte J, Maisetti O, Guillaume A, Skalli W, Portero P. Neck strength and EMG activity in fighter pilots with episodic neck pain. *Aviat Space Environ Med.* 2008; 79(10):947–952.
31. Liberati A, Altman DG, Tetzlaff J, Gøtzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med.* 2009; 6(7):e1000100.
32. Lombard M, Snyder-Duch J, Bracken CC. Content analysis in mass communication: assessment and reporting of intercoder reliability. *Hum Commun Res.* 2002; 28(4):587–604.
33. McCarthy MJH, Grevitt MP, Silcocks P, Hobbs G. The reliability of the Vernon and Mior neck disability index, and its validity compared with the short form-36 health survey questionnaire. *Eur Spine J.* 2007; 16(12):2111–2117.
34. McHugh ML. Interrater reliability: the Kappa statistic. *Biochem Med (Zagreb).* 2012; 22(3):276–282.
35. Moon BJ, Choi KH, Yun C, Ha Y. Cross-sectional study of neck pain and cervical sagittal alignment in air force pilots. *Aerosp Med Hum Perform.* 2015; 86(5):445–51.
36. Muller A, Robb O, Ngo A, Cronin C. Developing Air Force's Combat Aircrew: Department of Defence. Canberra: Australian National Audit Office; 2004. Report No.: 47.
37. Netto K, Hampson G, Oppermann B, Carstairs G, Aisbett B. Management of neck pain in Royal Australian Air Force fast jet aircrew. *Mil Med.* 2011; 176(1):106–109.
38. Newman DG. *Flying fast jets: human factors and performance limitations.* Boca Raton (FL): CRC Press; 2017:50–61.
39. Newman DG. +GZ-induced neck injuries in Royal Australian Air Force fighter pilots. *Aviat Space Environ Med.* 1997; 68(6):520–524.
40. Paganini-Hill A, Hsu G, Chao A, Ross RK. Comparison of early and late respondents to a postal health survey questionnaire. *Epidemiology.* 1993; 4(4):375–379.
41. Riechert M, Biesenbender S, Quix C. Developing and standardising definitions for research information: framework and methods of successful process documentation. *Procedia Comput Sci.* 2017; 106:260–267.
42. Rintala H, Häkkinen A, Siitonen S, Kyröläinen H. Relationships between physical fitness, demands of flight duty, and musculoskeletal symptoms among military pilots. *Mil Med.* 2015; 180(12):1233–1238.
43. Science and Technology Organization/North Atlantic Treaty Organization. HFM-252 Research Task Group on Aircrew Neck Pain. STO.NATO.INT. 2014 [Accessed Aug. 2019]. Available from: <https://www.sto.nato.int/sitepages/newsitem.aspx?id=2402>.
44. Sepucha KR, Matlock DD, Wills CE, Ropka M, Joseph-Williams N, et al. "It's valid and reliable" is not enough: critical appraisal of reporting of measures in trials evaluating patient decision aids. *Med Decis Making.* 2014; 34(5):560–566.
45. Shiri R, Frilander H, Sainio M, Karvala K, Sovelius R, et al. Cervical and lumbar pain and radiological degeneration among fighter pilots: a systematic review and meta-analysis. *Occup Environ Med.* 2015; 72(2):145–150.
46. Smith B, Chu LK, Smith TC, Amoroso PJ, Boyko EJ, et al. Challenges of self-reported medical conditions and electronic medical records among members of a large military cohort. *BMC Med Res Methodol.* 2008; 8(1):37.
47. Soeters J, Shields PM, Rietjens S. *Routledge handbook of research methods in military studies.* London (UK): Taylor & Francis; 2014:220–224.
48. Sovelius R, Oksa J, Rintala H, Huhtala H, Ylinen J, Siitonen S. Trampoline exercise vs. strength training to reduce neck strain in fighter pilots. *Aviat Space Environ Med.* 2006; 77(1):20–25.
49. Thoolen SJ, van den Oord MH. Modern air combat developments and their influence on neck and back pain in F-16 pilots. *Aerosp Med Hum Perform.* 2015; 86(11):936–941.
50. Timpka T, Alonso JM, Jacobsson J, Junge A, Branco P, et al. Injury and illness definitions and data collection procedures for use in epidemiological studies in Athletics (track and field): consensus statement. *Br J Sports Med.* 2014; 48(7):483–490.
51. Tucker B, Netto K, Hampson G, Oppermann B, Aisbett B. Predicting neck pain in Royal Australian Air Force fighter pilots. *Mil Med.* 2012; 177(4):444–450.
52. Vanderbeek RD. Period prevalence of acute neck injury in U.S. Air Force pilots exposed to high G forces. *Aviat Space Environ Med.* 1988; 59(12):1176–1180.
53. Van Loon AJ, Tijhuis M, Picavet HS, Surtees PG, Ormel J. Survey non-response in the Netherlands: effects on prevalence estimates and associations. *Ann Epidemiol.* 2003; 13(2):105–110.
54. VanValkenburg KR, Thompson A. Musculoskeletal pain in high-G aircraft training programs: a survey of student and instructor pilots. Wright-Patterson AFB (OH): USAF School of Aerospace Medicine; 2016. Technical Report AFRL-SA-WP-SR-2016-001. [Accessed 1 Aug. 2019.] Available from: <https://apps.dtic.mil/dtic/tr/fulltext/u2/1017966.pdf>.
55. Verde P, Trivelloni P, Angelino G, Morgagni F, Tomao E. Neck pain in F-16 vs. Typhoon fighter pilots. *Aerosp Med Hum Perform.* 2015; 86(4):402–406.
56. Wagstaff AS, Jahr KI, Rodskier S. +Gz-induced spinal symptoms in fighter pilots: operational and individual associated factors. *Aviat Space Environ Med.* 2012; 83(11):1092–1096.
57. Whitley BE, Kite ME. *Principles of Research in Behavioral Science*, 3rd ed. New York (NY): Taylor & Francis; 2012:149–150.
58. Wongpakaran N, Wongpakaran T, Wedding D, Gwet KL. A comparison of Cohen's Kappa and Gwet's AC1 when calculating inter-rater reliability coefficients: a study conducted with personality disorder samples. *BMC Med Res Methodol.* 2013; 13(1):61.