

Developing, Implementing, and Applying Novel Techniques During Systematic Reviews of Primary Space Medicine Data

Andrew Winnard; Nick Caplan; Claire Bruce-Martin; Patrick Swain; Rochelle Velho; Roberto Meroni; Virginia Wotring; Volker Damann; Tobias Weber; Simon Evetts; Joanthan Laws

- BACKGROUND:** The Aerospace Medicine Systematic Review Group was set up in 2016 to facilitate high quality and transparent synthesis of primary data to enable evidence-based practice. The group identified many research methods specific to space medicine that need consideration for systematic review methods. The group has developed space medicine specific methods to address this and trialed usage of these methods across seven published systematic reviews. This paper outlines evolution of space medicine synthesis methods and discussion of their initial application.
- METHODS:** Space medicine systematic review guidance has been developed for protocol planning, quantitative and qualitative synthesis, sourcing gray data, and assessing quality and transferability of space medicine human spaceflight simulation study environments.
- RESULTS:** Decision algorithms for guidance and tool usage were created based on usage. Six reviews used quantitative methods in which no meta-analyses were possible due to lack of controlled trials or reporting issues. All reviews scored the quality and transferability of space simulation environments. One review was qualitative. Several research gaps were identified.
- CONCLUSION:** Successful use of the developed methods demonstrates usability and initial validity. The current space medicine evidence base resulting in no meta-analyses being possible shows the need for standardized guidance on how to synthesize data in this field. It also provides evidence to call for increasing use of controlled trials, standardizing outcome measures, and improving minimum reporting standards. Space medicine is a unique field of medical research that requires specific systematic review methods.
- KEYWORDS:** space medicine, systematic reviews, methodological guidance.

Winnard A, Caplan N, Bruce-Martin C, Swain P, Velho R, Meroni R, Wotring V, Damann V, Weber T, Evetts S, Laws J. *Developing, implementing, and applying novel techniques during systematic reviews of primary space medicine data. Aerosp Med Hum Perform.* 2021; 92(8):681–688.

The Aerospace Medicine Systematic Review Group (AMSRG) was set up in 2016 to facilitate high quality and transparent synthesis of primary data to enable evidence-based medical practice. The benefits of the group were outlined in the May 2017 *Aerospace Medicine and Human Performance* journal President's Page as developing the knowledge base and improving the quality and value of research while highlighting research gaps to strengthen arguments for funding by demonstrating research needs. International gold standards for synthesizing traditional medical data are outlined generally in the Cochrane handbook⁷ and should be reported to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses

From the Aerospace Medicine and Rehabilitation Laboratory, Faculty of Health and Life Sciences, Northumbria University, Newcastle Upon Tyne, United Kingdom; the Department of Physiotherapy, LUNEX International University of Health, Differdange, Luxembourg; Human Performance in Space, International Space University, Strasbourg, France; the Space Medicine Team, ESA HRE-OM, European Astronaut Centre (EAC) and KBR GmbH, Cologne, Germany; and Blue Abyss, Liverpool, United Kingdom.

This manuscript was received for review in October 2020. It was accepted for publication in March 2021.

Address correspondence to: Andrew J. Winnard, Ph.D., Lecturer in Clinical/ Musculoskeletal Biomechanics, Northumbria University, Northumberland Bldg., Newcastle Upon Tyne NE1 8ST, United Kingdom; ajwinnard@gmail.com.

Reprint and copyright © by the Aerospace Medical Association, Alexandria, VA.

DOI: <https://doi.org/10.3357/AMHP.5803.2021>

Table I. AMSRG Reviews and Methods Used.

REVIEW	AIM	AMSRG METHODS USED
Valayer <i>et al.</i> ^{27†}	Evaluate if caloric restriction and dietary fasting can mitigate the adverse effects of ionizing radiation for deep space exploration.	Protocol & prescoping. Quantitative methods guide. Effect size analysis. Significance result reporting.
Sandal <i>et al.</i> ^{25†}	Evaluate the effectiveness of nutritional countermeasures as a standalone intervention to ameliorate musculoskeletal and cardiopulmonary deconditioning in gravitationally unloaded humans.	Protocol & prescoping. Bed rest quality score. Quantitative methods guide. Effect size analysis.
Laws <i>et al.</i> ¹³	Identify the technical constraints of the Orion multi-purpose crew vehicle or transferable spacecraft that impact on the capability of astronauts to exercise effectively.	Protocol & prescoping. Quality of nonempirical data sources score. Qualitative methods guide to conduct a thematic analysis.
Winnard <i>et al.</i> ^{32†}	Determine the time when humans exposed to simulated microgravity while not performing countermeasures reach a moderate or large effect on muscle health outcomes linked to if astronauts could have periods of no exercise on Moon or Mars missions.	Protocol & prescoping. Quantitative methods guide.* Bed rest quality score. Effect size analysis.
Konda <i>et al.</i> ¹⁰	Investigate exercise countermeasures for attenuating musculoskeletal deconditioning during long duration bed rest.	Protocol & prescoping. Quantitative methods guide.* Significance result reporting.
Winnard <i>et al.</i> ^{31†}	Assess interventions for counteracting changes and reducing injury risks to the lumbopelvic region during microgravity exposure	Protocol & prescoping.* Bed rest quality score.* Quantitative methods guide.* Effect size analysis.*
Richter <i>et al.</i> ^{24†}	Determine the human cardiopulmonary and biomechanical changes expected to occur in partial gravity to inform future Moon and Mars mission operations.	Protocol & prescoping.* Quality of partial gravity simulations tool.* Quantitative methods guide.* Effect size analysis.*

*Prepublication development version;

†EAC collaborative review.

(PRISMA) guidelines.¹⁷ Individual specialist topic areas tend to establish groups² to address their specific research intricacies and practices. The most thorough example of specialist groups are those within Cochrane; however, Cochrane groups require substantial recurring funding to set up and maintain databases of centrally quality scored trials, up to date methods, protocol registrations, and systematic review publication databases staffed by information specialists. The AMSRG is a good initial step toward high-quality regulation while funding is limited. All good quality systematic reviews should also conform to PRISMA guidelines. Aerospace is one of the final remaining fields of medicine to do so, a gap the AMSRG is addressing.

AMSRG has identified that space medicine has many research methods and limitations that are specific to the field and impact the systematic review process, such as small sample sizes, a lack of controlled trials, and common usage of ground-based spaceflight analogs. These aspects necessitate adaptation of systematic review synthesis methods and often require consideration regarding how findings from different research settings may safely transfer to the operational space medicine environment.¹⁸ To date, space medicine has relied on individual operational expert opinions or nonstandardized, evidence books¹⁹ that do not adhere to internationally recognized systematic review standards. In addition, the sample sizes of spaceflight and many analog studies are also small compared to those used in terrestrial clinical medicine, which can lead to more individual observations. This prevents standardized, transparent, repeatable, and easily updatable syntheses within which risk of bias, certainty, and transferability can be addressed. The

AMSRG has published seven systematic reviews as a group, five done in collaboration with the European Astronaut Centre on operationally driven topics (highlighted in **Table I**). During these reviews, space medicine specific methods and decision algorithms have been developed and trialed by the AMSRG. This paper provides a summary of these methods, trial usage, discussion for ongoing usage, and development from research, operational, and political perspectives.

METHODS

Protocol Planning and Prescoping

The AMSRG follows Cochrane guidance that requires initial planning and documenting methods decisions in a written protocol. Protocols state a clear question, scope, search strategy, inclusion criteria, and analysis decisions, such as subgroupings and statistical choices. Importantly, protocols record these decisions before results are available to reduce potential reviewer bias around methods decisions after synthesis results are available. The protocol stage also helps ensure a manageable scope likely to return the required data to run the statistical methods selected. While Cochrane requires a fully published protocol,³ the AMSRG strongly recommends a protocol and prescoping step,²⁸ in which decisions are made on search terms, scope, quality scoring, and synthesis methods and quick, prescoping searches are performed to check existence of relevant data. For a review to be considered systematic, the protocol and prescoping must result in a systematic search strategy using Boolean

TABLE 1 | Search strategy.

Search number	Term	Keywords in Boolean search format	Search mask
1	Partial gravity	"partial gravity" OR "fractional gravity" OR "reduced gravity" OR "lunar gravity" OR "moon gravity" OR "martian gravity" OR "mars gravity" OR "1/8th gravity" OR "1/8 G" OR "1/3rd gravity" OR "1/3 G" OR "low gravity" OR hypogravity OR "partial-gravity" OR "reduced-gravity" OR "Hypogravity" [Mesh;NoExp]	Title/ Abstract
2	Musculoskeletal	muscle* OR muscle OR bone* OR bone OR skeletal OR musculoskeletal OR "lean body mass" OR "body composition" OR osteo* OR osteo OR "musculo-skeletal" OR neuromusculoskeletal OR "Musculoskeletal System" [Mesh]	All Fields
3	Cardiopulmonary	cardio* OR cardio OR cardiac OR pulmona* OR pulmonary OR cardiopulmonary OR cardiovascular OR vascular* OR vascular OR respiratory OR respiration OR physiolog* OR physiological OR physiology OR heart* OR heart OR blood* OR blood OR capillarisation OR capillary OR myocard* OR myocard OR arterial OR venous OR orthostatic OR energetic* OR energetic OR energy OR metabolic OR OR "Cardiovascular System" [Mesh] OR "Blood" [Mesh] OR "Circulatory and Respiratory Physiological Phenomena" [Mesh]	All Fields
4	Mechanics	biomechanic* OR biomechanics OR mechanic* OR mechanic OR locomotion OR gait OR walk* OR walk OR run* OR run OR jump* OR jump OR landing OR "ground reaction forces" OR impact* OR impact OR "EMG" OR electromyo* OR electromyography OR "mechanical work" OR kinetics OR kinematics OR workload OR power OR "Movement" [Mesh] OR "Mechanics" [Mesh] OR "Mechanical Phenomena" [Mesh]	All Fields
5	Partial g simulations and methods	("body weight support" OR harness OR "alterG" OR "water immersion" OR "tilt table" OR "head-up tilt" OR "parabolic flight" OR "tail suspension" OR "supine suspension" OR "LBPP" OR "lower body positive pressure" OR "pressure suit" OR "subjects load device" OR centrifug* OR centrifugation OR "vertical treadmill" OR exoskeleton) AND gravity	All Fields
7	Combined search	1 AND (2 OR 3 OR 4 OR 5)	

Keywords were combined using the Boolean operators and grouped by main search terms. Medical Subject Headings (MeSH) as a comprehensive controlled vocabulary for the purpose of indexing journal articles and books in the life sciences were included in the search strategy. In the Pubmed advanced search builder either "Title/Abstract" or "All Fields" was used. The combined search allows to screen databases for various combinations of main search terms and their keywords.

Fig. 1. Example search strategy table from Richter et al. (2017).²⁴

logic and inclusion criteria that is detailed in the final publication alongside valid quality scoring and synthesis methods to enable the review to be transparent and repeatable. An example search strategy table to find studies detailing human biomechanical and cardiopulmonary changes due to partial gravity from a AMSRG and European Astronaut Centre (EAC) collaborative review is presented in **Fig. 1**.

As space medicine has many sources of data that sit outside common medical journal databases, the AMSRG provides a list of search locations to help locate gray literature, such as technical reports, for inclusion in reviews.³⁰ With regards to protocol planning, guides to help with both qualitative and quantitative methods have been made available. These guides have evolved over the years and have been modified by the AMSRG group.^{12,29}

Deciding Main Synthesis Method

To date, AMSRG has followed a decision algorithm that primarily recommends full quantitative meta-analysis and preference for reviews to be based on controlled clinical trials, where possible, following Cochrane quantitative methods. However, based on the initial review questions tackled, it is apparent that controlled trials, repeated studies, and use of standardized outcome measures across studies often do not exist or have been poorly reported, making Cochrane based meta-analyses not possible for some research questions. Therefore, alternate acceptable analyses were performed in a hierarchy of effect size

analysis, qualitative analyses^{20,26} (individually or combined with quantitative analysis within an integrated approach),⁹ and, finally, if no other method was possible and there was a justified reason for continuing, reporting of significance testing results from included papers.²⁷ In extreme cases, if no published data exists on a key topic, an empty review could be published to provide comprehensive gap analysis and be used to stimulate primary research. However, care should be taken to ensure an empty review is not solely due to an overly specific question.³⁴ The methods decision algorithm is presented in **Fig. 2**.

Effect size analysis converts all reported data to standardized units that can be reported in a single unit with a confidence interval to enable comparison across studies and outcomes to identify overall trends. The AMSRG recommends considering the use of effect size bias correction using the Hedges method that corrects for small sample sizes common in space medicine.¹¹ The AMSRG provides a spreadsheet to manage data extraction and calculates both basic and Hedges corrected effect sizes.³³ Results are still displayed on Forest plots, but effect size analysis does not complete the final meta-analysis step, so there is no overall synthesis statistic, heterogeneity step, or diamond to represent pooled effect on the Forest plots. The Cochrane handbook refers to this as "summarizing effect estimates."¹⁶

Qualitative methods are useful when data, technical reports, and discussions clearly exist in published materials on an important topic, but not within controlled trials. These methods




Fig. 2. Methods decision algorithm with the decisions running down the middle starting from the top. Dashed lines show alternate or supplementary decision options.

are also useful to combine with quantitative analysis to capture a more holistic multinodal dialogic explanation of a research topic;¹⁵ this is known as an integrated approach.^{4,21} The AMSRG qualitative guidance centers around thematic analysis, identifying common themes across the evidence base before structuring them into thematic maps that explore the relationships and potential hierarchies that exist within the data, and potentially integrating quantitative and qualitative data together to provide insights toward answering research questions.^{1,26}

Significance test reporting has not been formally supported by the AMSRG quantitative methods guides to date as it is limited compared to meta and effect analysis. However, as it has been used by a small number of initial reviews due to lack of effect size data it has now been added to the decision algorithm. The method involves reporting the *P*-values of studies when there is no other data available to calculate effect sizes, but there are strong justifications for still acknowledging the results of some studies.

Developing Options for Quality Scoring

Risk of bias has been assessed using Cochrane risk of bias tools^{6,8} for any controlled or within participant trials. However, it is very specific to randomized controlled trials and so has often not been fully applicable to space medicine studies. In addition, the Cochrane tool is not suited to scoring many gray literature sources such as technical reports. Several tools have

been developed and trialed for scoring quality, usage, and transferability of human spaceflight terrestrial based analogs. A bed rest quality score reports both greater quality and transferability as a higher score³¹ on an 8-point scale that considers various factors considered important to the quality of bed rest as an astronaut microgravity simulation. A rank has been compiled that lists how well various ground simulations are likely to accurately model astronaut biomechanical, cardiovascular, and metabolic changes. This gives an indication of how safe it is to transfer findings from the models to astronaut medicine.²³ For reviews including gray literature, there is a tool for scoring the quality of nonempirical data sources such as technical documents. This tool is based on greater scoring of documents if they are well sourced, clearly written, and based upon previous research as opposed to documents lacking citations and not having clearly explained methods for how findings or conclusions were reached.¹⁴ An algorithm to help decide which tools might be used for various types of reviews is shown in Fig. 3.

RESULTS

As of August 2020, seven space medicine systematic reviews had been completed and published in peer reviewed scientific journals (see Table I). Six of the reviews used quantitative




Fig. 3. Decision algorithm for considering use of various tools.

analysis and one used qualitative. All the reviews conformed to PRISMA standards and followed Cochrane guidelines wherever they were not using AMSRG specific methods, guides, and tools based on the decision algorithms. AMSRG specific methods were used in every instance to assess quality of space medicine specific data, including space simulation analog research environments and technical reports. All of the reviews were human space medicine topics. In all quantitative reviews, a full meta-analysis was not possible as the outcomes were highly heterogeneous across studies and there was poor reporting of space medicine studies. However, all quantitative studies were able to perform effect size analysis and only two^{10,27} had to rely on significance result reporting. One review¹³ used qualitative methods to present a thematic analysis of technical documents from gray literature sources required to answer the technical nature of the research question. Three reviews^{25,31,32} included bed rest spaceflight simulation studies and reported bed rest quality scores. One review²⁴ used the quality of partial gravity

simulations to list how well various ground simulations are likely to accurately model astronaut biomechanical, cardiovascular, and metabolic changes. A summary of all seven reviews, the research questions each posed, and the AMSRG methods used within the methods is presented in Table I.

Several gaps have been identified from the published reviews. Gaps are explained within each review and this is recommended by the AMSRG for all reviews. To provide an easily accessible summary overview of the gaps, they are also listed on the AMSRG gap analysis web page (<http://aerospacemed.rehab/gap-analysis>). A very brief summary of the gaps listed on the webpage as of August 2020 is as follows:

- Space agencies are advised to make information available about medically relevant constraints of spacecraft as accessible publications rather than solely in gray literature.
- Additional systematic reviews should carry on from the initial muscle review to establish the time for onset of

musculoskeletal and cardiovascular effects without counter-measures for potential no-exercise periods during space missions.

- More primary data are needed on expected muscle changes in microgravity when not taking exercise countermeasures.
- Primary research data linked to muscle changes in spaceflight need to standardize outcome measures used and consider including patient reported measures within the standardized list.
- More studies are required to investigate the countermeasures for minimizing risk to the lumbopelvic region during spaceflight.
- More data on the expected human physiological effects of various g-loading environments are needed to inform medical operations for lunar and Martian missions.

DISCUSSION

Seven AMSRG methods specific to space medicine systematic reviews which cover both quantitative and qualitative analysis have been developed to address the unique limitations, research gaps, and challenges of space medicine research. Decision algorithms to guide researchers in which AMSRG methods to use have been trialed. Having AMSRG as a central group to standardize and publish review methods, along with a summary of any identified gaps, is beneficial to managing information in a single place, providing sector oversight, and improving research quality in space medicine.

Developing an Algorithm to Decide the Best Review

Method to Use

Space medical research presents challenges that have prevented the use of meta-analysis in the AMSRG's current reviews. These challenges include the use of different spaceflight simulation models, a lack of controlled clinical trials, many heterogeneous outcome measures, a lack of standardization, and, in some cases, poor reporting of study data. The identification of these challenges by AMSRG and the development or adaptation of methods to address them is evidence of the need for a centralized group to provide this guidance and gap analysis for the sector. The AMSRG decision algorithm, supported by the Cochrane handbook, prioritizes analyses in the order of: meta-analysis; effect size analysis; and then significance test reporting. The algorithm provides the option of considering qualitative analysis at all levels, either to supplement quantitative data in an integrative approach or as an alternative when data is available but in a format that does not support quantitative methods. As significance testing is now a formal accepted step in AMSRG methods, the AMSRG quantitative guidance will be updated to include additional significance testing options, such as combining *P*-values and vote counting, to bring the AMSRG guidance fully in line with all valid non-meta-analysis options suggested by Cochrane. It should be noted that there are limitations to the non-meta-analysis methods

detailed in the Cochrane handbook, and so caution must be taken when using them. For example, care must still be taken when using effect size and/or *P*-value-based analyses to recognize the original units of included measures and not make unreasonable comparisons such as comparing wildly differing outcomes. In addition, the reasons why space medicine reviews are forced to use alternate methods should be identified and addressed and the AMSRG has recommended increasing performing of controlled clinical trials, determining and then using standardized outcome measures relevant to space operations, and ensuring reporting standards of space medicine research supports systematic reviews. Reporting all experimental group means, standard deviations, and group sample sizes is required for meta-analysis and should be set as the minimum journal publishing requirements in addition to reporting any statistical results such as *P*-values. Furthermore, basic data on medical requirements/constraints of human spaceflight environments such as the internal volume of spacecraft that is available for operational essential activities, including exercise countermeasures, should be made easily available and accessible to the research community.

Quality Scoring Tools for Space Medicine

The wide variety of ground-based space simulations used in research need to be assessed for risk of bias, quality, and transferability of studies using specifically tailored methods as opposed to generic quality tools. To address this for bed rest, the most commonly encountered simulation within the published reviews to date, a specific quality scoring tool was developed. This was developed by communicating with a team of experts to establish and then agree to the final criteria and was done as part of one of the first completed reviews.³¹ This has since been supplemented with an AMSRG ranking system that indicates how well parabolic flight, bed rest, isolation, and suspension studies' findings can be transferred to actual astronaut settings during spaceflight.

However, additional detailed quality scoring tools for each environment would be beneficial to develop. To date, where spaceflight data did not exist, AMSRG reviews have remained within the scope of human simulation research only, as there were concerns that animal models might be too severely limited for transferability. In some cases, nonempirical sources may also provide useful insights at a human level and the AMSRG has provided a quality assessment tool for such documents. The use of the AMSRG developed tools in successfully published reviews, several of which have been done in collaboration with operational space agency medical staff from the European Space Agency, shows an initial level of validity of the tools. Going forward, it would be useful to also test both the inter- and intrarater reliability of the tools. Both reliabilities are important as best practice is to use agreement of multiple reviewers when scoring papers. While the methods for scoring space medicine studies reported in this document consider quality and transferability, it has been mostly possible to use Cochrane's risk of bias tool for assessing bias. Where study designs made the Cochrane risk of bias

inappropriate, alternative validated tools were available such as the Physiotherapy Evidence Database scale²² and Quality in Prognostic Studies.⁵ It is likely that review teams will be able to use existing tools such as these to score bias and use AMSRG tools to further assess the quality and transferability of the various ground-based simulations.

In conclusion, space medicine is a unique field of medical research that requires specific systematic review methods to be developed to enable safe, transparent, reproducible synthesis of primary data to develop a robust evidence base that underpins space medical operations. After performing seven systematic reviews in aerospace medicine, adopting traditional systematic review tools by the Cochrane group has been challenging and has required modification to capture the full breadth of primary sources available in aerospace medicine. The AMSRG group has built and will continue to build on the relevant foundations required to curate a central repository of educational resources which are required to perform systematic reviews in aerospace medicine using evidence-based methodology without compromising scientific rigor.

The limitations of the methods covered here are in the developmental nature of them. As already mentioned, the tools have been developed and then trialed within the initial reviews to establish them and test their validity and usability in the field. It would be useful to also establish the reliability of the tools that involve author scoring of studies.

ACKNOWLEDGEMENTS

Financial Disclosure Statement: The authors have no competing interests to declare.

Authors and Affiliations: Andrew J. Winnard, Ph.D., M.Sc., Nick Caplan, Ph.D., Pg.Cert., Clair Bruce-Martin, M.Sc., PGCAPI, Patrick Swain, M.Sc., B.Sc., Rochelle Velho, M.B.Ch.B., M.P.H., and Jonathan M. Laws, M.Res., B.Sc., Aerospace Medicine and Rehabilitation Laboratory, Faculty of Health and Life Sciences, Northumbria University, Newcastle Upon Tyne, United Kingdom; Roberto Meroni, Ph.D., M.Sc., Department of Physiotherapy, LUNEX International University of Health, Differdange, Luxembourg; Virginia Wotring, Ph.D., B.S., and Volker Damann, M.D., Human Performance in Space, International Space University, Strasbourg, France; Tobias Weber, Ph.D., B.Sc., Space Medicine Team, ESA HRE-OM, European Astronaut Centre (EAC) and KBR GmbH, Cologne, Germany; and Simon Evetts, Ph.D., M.Sc., Blue Abyss, Liverpool, United Kingdom.

REFERENCES

1. Castleberry A, Nolen A. Thematic analysis of qualitative research data: is it as easy as it sounds? *Curr Pharm Teach Learn*. 2018; 10(6):807–815. Erratum in: *Curr Pharm Teach Learn*. 2021; 13(2):192.
2. Cochrane. Cochrane Review groups and networks. 2020; [Accessed 06/10/2020]. Available from: <https://www.cochranelibrary.com/about/cochrane-review-groups>.
3. Cumpston M, Chandler J. Chapter II: planning a Cochrane Review. In: Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, et al., editors. *Cochrane Handbook for Systematic Reviews of Interventions*, 2nd ed. Chichester (UK): John Wiley & Sons; 2019:6.
4. Harden A, Thomas J. Mixed methods and systematic reviews: examples and emerging issues. In: Tashakkori A, Teddlie C, editors. *Sage handbook of mixed methods in social & behavioral research*. London: Sage; 2010; 2: 749–774.
5. Hayden JA, van der Windt DA, Cartwright JL, Côté P, Bombardier C. Assessing bias in studies of prognostic factors. *Ann Intern Med*. 2013; 158(4):280–286.
6. Higgins J, Altman D, Sterne J; Cochrane Statistical Methods Group and the Cochrane Bias Methods Group. Chapter 8: Assessing risk of bias in included studies. In: Higgins JP, Green S, editors. *Cochrane handbook for systematic reviews of interventions*, vs. 5.1.0. London: The Cochrane Collaboration; 2011.
7. Higgins JP, Savović J, Page MJ, Elbers RG, Sterne JA. Assessing risk of bias in a randomized trial. In: Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, et al., editors. *Cochrane Handbook for Systematic Reviews of Interventions*, 2nd ed. Chichester (UK): John Wiley & Sons; 2019:205–228.
8. Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, et al. *Cochrane Handbook for Systematic Reviews of Interventions*, version 6.0. 2019; [Accessed 23/03/2020]. Available from: <https://training.cochrane.org/handbook>.
9. Hong QN, Pluye P. A conceptual framework for critical appraisal in systematic mixed studies reviews. *J Mixed Methods Res*. 2019; 13(4):446–460.
10. Konda NN, Karri RS, Winnard A, Nasser M, Evetts S, et al. A comparison of exercise interventions from bed rest studies for the prevention of musculoskeletal loss. *NPJ Microgravity*. 2019; 5(1):12.
11. Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Front Psychol*. 2013; 4:863.
12. Laws J, Bruce-Martin C, Winnard A. Aerospace Medicine Systematic Review Group qualitative methods guide for space medicine focused systematic reviews. 2020; [Accessed 23/03/2020]. Available from: https://www.researchgate.net/publication/339911374_Aerospace_Medicine_Systematic_Review_Group_Qualitative_Methods_Guide_for_Space_Medicine_Focused_Systematic.
13. Laws J, Caplan N, Bruce C, McGrogan C, Lindsay K, et al. Systematic review of the technical and physiological constraints of the Orion multi-purpose crew vehicle that affect the capability of astronauts to exercise effectively during spaceflight. *Acta Astronaut*. 2020; 170:665–677.
14. Laws JM, Winnard A. Tool for scoring the quality of non-empirical data sources. E.G: Technical Reports 2019. [Accessed May 17, 2021]. Available from: https://www.researchgate.net/publication/331385312_Tool_for_Scoring_the_Quality_of_Non-Empirical_Data_Sources-EG_Technical_Reports.
15. Mason J. Mixing methods in a qualitatively driven way. *Qual Res*. 2006; 6(1):9–25.
16. McKenzie JE, Brennan SE. Synthesizing and presenting findings using other methods. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, et al., editors. *Cochrane Handbook for Systematic Reviews of Interventions*. London: The Cochrane Collaboration; 2019:321–347.
17. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009; 6(7):e1000097.
18. Morey-Holton ER. Ground-based models for studying adaptation to altered gravity. Moffet Field (CA): NASA Ames Research Centre Report; 2000.
19. NASA. Evidence book overview. 2020; [Accessed 2020 06/10/2020]. Available from <https://humanresearchroadmap.nasa.gov/evidence/#overview>.
20. Noyes J, Booth A, Cargo M, Flemming K, Harden A, et al. Qualitative evidence. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, et al., editors. *Cochrane Handbook for Systematic Reviews of Interventions*. London: The Cochrane Collaboration; 2019:525–545.
21. Pearson A, White H, Bath-Hextall F, Salmond S, Apostolo J, Kirkpatrick P. A mixed-methods approach to systematic reviews. *Int J Evid-Based Healthc*. 2015; 13(3):121–131.
22. PEDro. PEDro scale, the George Institute for Global Health. 1999; [Accessed 08/08/2015]. Available from: www.pedro.org.au.
23. Richter C, Braunstein B, Weber T. Tool for assessing quality of partial gravity simulation models. 2017; [Accessed 06/10/20]. Available from: https://www.researchgate.net/publication/319464193_Tool_for_Assessing_Quality_of_Partial_Gravity_Simulation_Models.

24. Richter C, Braunstein B, Winnard A, Nasser M, Weber T. Human biomechanical and cardiopulmonary responses to partial gravity—a systematic review. *Front Physiol.* 2017; 8:583.
25. Sandal PH, Kim D, Fiebig L, Winnard A, Caplan N, et al. Effectiveness of nutritional countermeasures in microgravity and its ground-based analogues to ameliorate musculoskeletal and cardiopulmonary deconditioning—a systematic review. *PLoS One.* 2020; 15(6):e0234412.
26. Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol.* 2008; 8(1):45.
27. Valayer S, Kim D, Fogtman A, Straube U, Winnard A, et al. The potential of fasting and caloric restriction to mitigate radiation damage—a systematic review. *Front Nutr.* 2020; 7:584543.
28. Winnard A. AMSRG protocol template. 2017; [Accessed 06/10/2020]. Available from: https://www.researchgate.net/publication/318541310_AMSRG_Protocol_Template.
29. Winnard A, Bruce-Martin C, Laws J. Aerospace Medicine Systematic Review Group Quantitative Methods Guide for Space Medicine Focused Systematic Reviews. 2020; [Accessed 23/03/2020]. Available from: https://www.researchgate.net/publication/339788931_Aerospace_Medicine_Systematic_Review_Group_Quantitative_Methods_Guide_For_Space_Medicine_Focussed_Systematic_Reviews.
30. Winnard A, Laws J, Swain P. AMSRG list of potential data sources. 2020; [Accessed 23/03/2020]. Available from: https://www.researchgate.net/publication/338774503_AMSRG_List_of_Potential_Data_Sources.
31. Winnard A, Nasser M, Debuse D, Stokes M, Evetts S, et al. Systematic review of countermeasures to minimise physiological changes and risk of injury to the lumbopelvic area following long-term microgravity. *Musculoskelet Sci Pract.* 2017; 27(Suppl. 1):S5–S14.
32. Winnard A, Scott J, Waters N, Vance M, Caplan N. Effect of time on human muscle outcomes during simulated microgravity exposure without countermeasures—systematic review. *Front Physiol.* 2019; 10:1046.
33. Winnard A, Waters N. AMSRG study information and data extraction spreadsheet. 2018; [Accessed May 17, 2021]. Available from: https://www.researchgate.net/publication/326958082_AMSRG_study_information_and_data_extraction_spreadsheet/citation/download.
34. Yaffe J, Montgomery P, Hopewell S, Shepard LD. Empty reviews: a description and consideration of Cochrane systematic reviews with no included studies. *PLoS One.* 2012; 7(5):e36626.